

Spin-Wave Theory of the Spiral Phase of the t - J Model

N.I.Karchev

Department of Physics, University of Sofia, 1126 Sofia, Bulgaria

T.S.Hristov

*Space Research Institute, Bulgarian Academy of Sciences, 1000 Sofia,
Bulgaria*

Abstract

A graded Holstein-Primakoff realization of the $SU(2/1)$ algebra is proposed. A spin-wave theory with a condition that the sublattice magnetization is zero is discussed. It is shown that the generalized spin-wave theory is appropriate to investigate the long-range spiral (incommensurate) phase of the $t - J$ model at $T = 0$. The spin-spin correlator is calculated.

PACS Numbers: 75.10.Jm, 74.20.-z, 74.65.+n

I. Introduction

Among the many electronic models which are being currently studied in the context of high- T_c superconductivity, the two dimensional $t - J$ model is the simplest one and it is rather general to describe the basic physics of the new superconductor¹. It is an effective model for the large U Hubbard model with $J = \frac{4t^2}{U}$, and there are arguments that one can derive it from a multiband Hubbard model^{2,3}.

At half-filling the $t - J$ model reduces to the Heisenberg model. This is a welcome result, since, the methods developed for the Heisenberg model can be extended and applied to investigate the properties of the $t - J$ model, atleast at nearly half-filling. Schwinger boson slave-fermion mean field theory was used and an antiferromagnetic ordered Neél state at half-filling was found⁴. This state evolves into long-range spiral (incommensurate) antiferromagnetic states at large doping. Quantitatively, this is described by the dependence of the spirality angle Q on the doping⁵. Originally, the spiral phase of the $t - J$ model was obtained by Shraiman and Siggia who used a somewhat different approach⁶.

Recently, Takahashi has formulated a modified spin-wave theory of Heisenberg (anti)ferromagnets^{7,8}. He has supplemented the usual spin-wave theory with the constraint that the magnetization of each site is zero. This enforces the condition that the total number, on average, of spin waves per site is S , and that the sublattice rotational symmetry is not broken. In one dimension the modified spin wave theory yields excellent agreement with Bethe ansatz results for spin $S = 1/2$. Takahashi's results are in quantitative agreement with the Schwinger boson mean field theory⁴ and with the renormalization group theory⁹.

In this paper we formulate, along the same line, a generalized spin wave theory appropriate to investigate the spiral phase of $t - J$ model. Some results of Ref. 10 are used.

In Section II the canonical bose- and fermi- operators are used to realize

the $SU(2/1)$ algebra. The graded Holstein-Primakoff representation is written in a local frame. The effective hamiltonian is obtained in leading order of S^{-1} , as a function of the angles θ_i which determine the local frame. To preserve the sublattice rotational symmetry we impose two additional conditions. To enforce them, we introduce two terms in the hamiltonian with chemical potentials λ and ζ . For the special values of $\theta_i = k\pi$, ζ is equal to zero and we reobtain the Takahashi's spin-wave theory. We express the angles θ_i by the spirality angle Q . To determine the last one, we calculate the energy of the system per site. The physical value of Q is that which minimizes the energy. It depends on the doping.

In Section III we calculate the large distance asymptotic of the spin-spin correlation function. At $T = 0$ we obtain long-range correlation which is characterized by the angle Q and the quantity m which is related to the spin-wave bose-condensate. At finite temperature, the correlation function exponentially falls when the separation between the two sites is large. At small doping the spin-spin correlator looks like in the theory of Heisenberg antiferromagnet. When the doping is large enough it looks like in the theory of Heisenberg ferromagnet.

Section IV is devoted to the concluding remarks.

II. Graded spin-wave theory

The $t - J$ model is defined by the Hamiltonian

$$h = -t \sum_{\langle i,j \rangle} [c_{i\sigma}^+ c_{j\sigma} + h.c.] + J \sum_{\langle i,j \rangle} \left(\vec{S}_i \cdot \vec{S}_j - \frac{1}{4} n_i n_j \right) - \mu \sum_i c_{i\sigma}^+ c_{j\sigma} \quad (1)$$

where $c_{i\sigma}^+(c_{j\sigma})$ are the fermi operators of the electron ($\sigma = 1, 2$) on site $i(j)$ of a 2D square lattice which act on states with no double occupancy on a lattice site $i(j)$, $\vec{S}_i$ are spin operators, and μ is the chemical potential. By $\langle i, j \rangle$ we denote the sum over the nearest neighbors.

The eight operators $c_{i\sigma}$, $c_{i\sigma}^+$, $S_i^\pm$, S_i^3 and n_i form a basis of the $SU(2/1)$ graded algebra. It is most convenient to rewrite them in terms of the Hubbard

operators X^{ab} ($a, b = 1, 2, 3$)

$$\begin{aligned}
c_{i\sigma} &= X_i^{3\sigma} & c_{i\sigma}^+ &= X_i^{\sigma 3} & \sigma &= 1, 2 \\
n_i &= X_i^{11} + X_i^{22} & S_i^3 &= \frac{1}{2} (X_i^{11} - X_i^{22}) \\
S_i^+ &= X_i^{12} & S_i^- &= X_i^{21}
\end{aligned} \tag{2}$$

They satisfy the following graded commutation rules

$$[X_i^{ab}, X_j^{cd}]_{\pm} = \delta_{ij} (\delta^{bc} X_i^{ad} \pm \delta^{ad} X_i^{cb}) \tag{3}$$

The graded algebra $SU(2/1)$ can be realized using a canonical bose a_i, a_i^+ and fermi ψ_i, ψ_i^+ operators

$$[a_i, a_j^+]_{-} = \delta_{ij}; \quad [\psi_i, \psi_j^+]_{+} = \delta_{ij} \tag{4}$$

$$\begin{aligned}
c_{i1} &= \psi_i^+ \left[(2S - a_i^+ a_i - \psi_i^+ \psi_i)^{1/2} \cos \frac{\theta_i}{2} + a_i \sin \frac{\theta_i}{2} \right] \\
c_{i1}^+ &= \left[(2S - a_i^+ a_i - \psi_i^+ \psi_i)^{1/2} \cos \frac{\theta_i}{2} + a_i^+ \sin \frac{\theta_i}{2} \right] \psi_i \\
c_{i2} &= \psi_i^+ \left[- (2S - a_i^+ a_i - \psi_i^+ \psi_i)^{1/2} \sin \frac{\theta_i}{2} + a_i \cos \frac{\theta_i}{2} \right] \\
c_{i2}^+ &= \left[- (2S - a_i^+ a_i - \psi_i^+ \psi_i)^{1/2} \sin \frac{\theta_i}{2} + a_i^+ \cos \frac{\theta_i}{2} \right] \psi_i \\
n_i &= 2S - \psi_i^+ \psi_i \\
S_i^3 &= \frac{1}{2} \sin \theta_i \left[(2S - a_i^+ a_i - \psi_i^+ \psi_i)^{1/2} a_i + a_i^+ (2S - a_i^+ a_i - \psi_i^+ \psi_i)^{1/2} \right] \\
&\quad + \cos \theta_i \left(S - a_i^+ a_i - \frac{1}{2} \psi_i^+ \psi_i \right) \\
S_i^+ &= \cos^2 \frac{\theta_i}{2} (2S - a_i^+ a_i - \psi_i^+ \psi_i)^{1/2} a_i - \sin^2 \frac{\theta_i}{2} a_i^+ (2S - a_i^+ a_i - \psi_i^+ \psi_i)^{1/2} \\
&\quad - \sin \theta_i \left(S - a_i^+ a_i - \frac{1}{2} \psi_i^+ \psi_i \right)
\end{aligned} \tag{5}$$

$$S_i^- = \cos^2 \frac{\theta_i}{2} a_i^+ (2S - a_i^+ a_i - \psi_i^+ \psi_i)^{1/2} - \sin^2 \frac{\theta_i}{2} (2S - a_i^+ a_i - \psi_i^+ \psi_i)^{1/2} a_i \\ - \sin \theta_i \left(S - a_i^+ a_i - \frac{1}{2} \psi_i^+ \psi_i \right)$$

where θ_i are arbitrary angles which run over intervals with length π . When $\theta_i = 0, \pi, 2\pi \dots$ the Eqs. (5) reduce to the graded Holstein-Primakoff representation¹¹. It is easy to prove that so defined operators satisfy the algebra Eqs.(2,3). In particular, the spin operators satisfy the SU(2) algebra and S parametrizes the representation of the SU(2/1) algebra. At half-filling S is exactly the spin of the system. The physical relevant case is $S = \frac{1}{2}$.

In spin-wave approximation one must replace $(2S - a_i^+ a_i - \psi_i^+ \psi_i)^{1/2}$ by $(2S)^{1/2} [1 - \frac{1}{4S} (a_i^+ a_i + \psi_i^+ \psi_i)]$ in Eqs. (5). Then, in leading order of S^{-1} , the Hamiltonian Eq. (1) takes the form

$$h = h_{ce} + h_q \quad (6)$$

where

$$h_{ce} = JS^2 \sum_{\langle i,j \rangle} \cos(\theta_i - \theta_j) \quad (7)$$

and

$$h_q = -JS \sum_{\langle i,j \rangle} \cos(\theta_i - \theta_j) (a_i^+ a_i + a_j^+ a_j) + \\ + \frac{JS}{2} \sum_{\langle i,j \rangle} [\cos(\theta_i - \theta_j) + 1] (a_i^+ a_j + a_j^+ a_i) \\ + \frac{JS}{2} \sum_{\langle i,j \rangle} [\cos(\theta_i - \theta_j) - 1] (a_i a_j + a_i^+ a_j^+) \\ - \frac{JS}{2} \sum_{\langle i,j \rangle} [\cos(\theta_i - \theta_j) - 1] (\psi_i^+ \psi_i + \psi_j^+ \psi_j) \\ + 2St \sum_{\langle i,j \rangle} \cos \frac{\theta_i - \theta_j}{2} (\psi_i^+ \psi_j + \psi_j^+ \psi_i) - \mu \sum_i (2S - \psi_i^+ \psi_i)$$

Farther on, we shall use special values for the angles θ_i

$$\theta_i = \vec{Q} \cdot \vec{r}_i; \quad Q_x = Q_y = Q \quad (9)$$

where $\pi \leq Q \leq 2\pi$ and $\vec{r}_i = (r_i^x, r_i^y)$ are the coordinates of the lattice site i .

The Eqs. (6-8) yield the conventional spin wave theory. To preserve the sublattice rotational symmetry one must impose, by hand, an additional condition. Here it is

$$\langle \vec{n}_i \cdot \vec{S}_i \rangle = 0 \quad (10)$$

where $\vec{n}_i$ is a unit vector which fixes the local frame we have chosen in Eqs.(5).

$$\vec{n}_i = (-\sin \theta_i, 0, \cos \theta_i) \quad (11)$$

When $\theta_i = 2k\pi$ the Eq.(10) is the condition imposed by Takahashi for ferromagnets⁷. When $\theta_i = k\pi$ Eq. (10) coincides with the condition discussed in spin-wave theory of antiferromagnets^{8,12}

Making use of Eq. (11) and the representation for the spin operators Eqs. (5), one can rewrite the condition Eq. (10) in the form

$$\langle a_i^+ a_i \rangle + \frac{1}{2} \langle \psi_i^+ \psi_i \rangle = S \quad (12)$$

To enforce the constraint Eq. (12) we introduce a new term in the Hamiltonian

$$h \rightarrow h - \lambda \sum_i (2a_i^+ a_i + \psi_i^+ \psi_i - 2S) \quad (13)$$

with Lagrange multiplier to be determined by Eq. (12).

It is not difficult to check, that Eq. (12) has a solution just for $Q = \pi, 2\pi$. This is easy to be understood because the identities

$$\langle a_i a_i \rangle = \langle a_i^+ a_i^+ \rangle = 0 \quad (14)$$

are satisfied only for these values of Q . To ensure the implementation of the Eq. (12) for arbitrary Q in the interval $[\pi, 2\pi]$, one must impose in addition the condition Eq. (14) also. We introduce in the Hamiltonian a term with Lagrange multiplier λ which enforces the condition Eq.(12) and a second term with Lagrange multiplier ζ which enforces the conditions Eq. (14). The full Hamiltonian in the momentum space reads

$$h = 2JS^2N \cos Q + 2SN(\lambda - \mu) \quad (15)$$

$$+ \sum_k \left\{ \epsilon_b(k) a_k^+ a_k + D_k [a_k a_{-k} + a_k^+ a_{-k}^+] + \epsilon_f(k) \psi_k^+ \psi_k \right\}$$

where

$$\begin{aligned} \epsilon_b(k) &= -2\lambda - 4SJ \cos Q + 2SJ(1 + \cos Q)\gamma_k \\ D_k &= SJ[\cos Q - 1]\gamma_k + \zeta \\ \epsilon_f(k) &= \mu - \lambda + 2SJ(1 - \cos Q) + 8St \cos \frac{1}{2}Q \gamma_k \\ \gamma_k &= \frac{1}{2}(\cos k_x + \cos k_y) \end{aligned} \quad (16)$$

N is the number of lattice sites, and $\vec{k}$ runs over the Brillouin zone. Throughout this paper we put the lattice spacing equal to one.

The Bose part of the Hamiltonian Eq. (15) is diagonalized by a Bogoliubov transformation. The result is

$$h_b = \sum_k [E_b(k) \alpha_k^+ \alpha_k + E_o(k)]$$

where

$$\begin{aligned} E_b(k) &= [\epsilon_b^2(k) - 4D_k^2]^{1/2} \\ E_o(k) &= \frac{1}{2} [E_b(k) - \epsilon_b(k)] \end{aligned} \quad (17)$$

The free energy of the system is given by the expression

$$\begin{aligned} F &= 2JS^2 \cos Q + 2S(\lambda - \mu) + \frac{1}{N} \sum_k E_o(k) \\ &+ \frac{1}{\beta N} \sum_k \ln(1 - e^{-\beta E_b(k)}) - \frac{1}{\beta N} \sum_k \ln(1 + e^{-\beta \epsilon_f(k)}) \end{aligned} \quad (18)$$

where $\beta = (k_B T)^{-1}$ is the inverse temperature. The three chemical potentials λ , ζ and μ are determined by the equations

$$\frac{\partial F}{\partial \lambda} = 0, \quad \frac{\partial F}{\partial \zeta} = 0, \quad \frac{\partial F}{\partial \mu} = 0$$

It is more convenient to write their linear combinations

$$\frac{\partial F}{\partial \lambda} + \frac{\partial F}{\partial \zeta} = 2S + 1 - \delta - \frac{1}{N} \sum_k \frac{\epsilon_b(k) + 2D_k}{E_b(k)} \text{cth} \frac{\beta}{2} E_b(k) = 0 \quad (19)$$

$$\frac{\partial F}{\partial \lambda} - \frac{\partial F}{\partial \zeta} = 2S + 1 - \delta - \frac{1}{N} \sum_k \frac{\epsilon_b(k) - 2D_k}{E_b(k)} \text{cth} \frac{\beta}{2} E_b(k) = 0 \quad (20)$$

$$\frac{1}{N} \sum_k \frac{1}{1 + e^{-\beta \epsilon_f(k)}} = \delta \quad (21)$$

where δ is the doping parameter.

We introduce new chemical potentials η_1, η_2 which are determined by the equalities

$$\begin{aligned} 2SJ\eta_1 &= -\lambda - \zeta - 2SJ \cos Q \\ 2SJ\eta_2 &= -\lambda + \zeta - 2SJ \cos Q \end{aligned} \quad (22)$$

Then

$$\begin{aligned} \epsilon_b(k) - 2D_k &= 4SJ(\eta_1 + \gamma_k) \\ \epsilon_b(k) + 2D_k &= 4SJ(\eta_2 + \cos Q \gamma_k) \\ E_b(k) &= 4SJ \sqrt{(\eta_1 + \gamma_k)(\eta_2 + \cos Q \gamma_k)} \end{aligned} \quad (23)$$

From Eqs.(19-23) it follows, that when $Q = \pi, 2\pi$, the solution is $\eta_1 = \eta_2$ or $\zeta = 0$. This means, that for these special values of Q the identifies Eq. (14) are satisfied without introducing a second term in the Hamiltonian and one reobtains the Takahashi's spin-wave theory.

The chemical potentials depend on the temperature and $\eta_1(T) \geq 1$; $\eta_2(T) \geq |\cos Q|$. Next, we analyze the low-temperature properties of Eqs. (19,20) $\eta_1(O)$ and $\eta_2(O)$ depend on Q and δ . For given doping δ we shall consider three cases.

When $\pi \leq Q \leq Q_1 \leq \frac{3}{2}\pi$, we obtain from Eqs. (19,20) (see Takahashi⁸)

$$\begin{aligned} 2S+1-\delta-1.394 |\cos Q|^{1/2} + \frac{1}{2\pi SJ\beta} \ln 2(4SJ\beta)^2 |\cos Q| (\eta_1 - 1) &= 0 \\ 2S+1-\delta-1.394 |\cos Q|^{-1/2} + \frac{1}{2\pi SJ\beta |\cos Q|} \ln 2(4SJ\beta)^2 (\eta_2 - |\cos Q|) &= 0 \end{aligned} \quad (24)$$

From Eq. (24) one gets

$$\eta_1 - 1 = \frac{1}{2(4SJ\beta)^2 |\cos Q|} e^{-4\pi SJm_1\beta}$$

$$\eta_2 - |\cos Q| = \frac{1}{2(4SJ\beta)^2} e^{-4\pi SJm_2|\cos Q|\beta} \quad (25)$$

where

$$\begin{aligned} 2m_1 &= 2S + 1 - \delta - 1.394 |\cos Q|^{1/2} \\ 2m_2 &= 2S + 1 - \delta - 1.394 |\cos Q|^{-1/2} \end{aligned} \quad (26)$$

m_1 increases and m_2 decreases as $|\cos Q|$ decreases, and $m_2 = 0$ for the upper bound of the interval

$$Q_{(1)} = \arccos \left[- \left(\frac{1.394}{2S + 1 - \delta} \right)^2 \right]$$

When $\frac{3\pi}{2} < Q_2 \leq Q \leq 2\pi$ one obtains from Eqs. (19,20)

$$\begin{aligned} 2S + 1 - \delta - (\cos Q)^{1/2} &= \\ &= -\frac{1}{2\pi SJ\beta} \ln 18SJ\Delta_1 \frac{(\Delta_1 + \Delta_2)^{1/2} + (\Delta_1 + 4\Delta_2)^{1/2}}{[2(\Delta_1 + \Delta_2)^{1/2} + (\Delta_1 + 4\Delta_2)^{1/2}]^2} \frac{\beta}{(\cos Q)^{1/2}} \\ 2S + 1 - \delta - (\cos Q)^{-1/2} &= \\ &= -\frac{1}{2\pi SJ \cos Q \beta} \ln 2SJ[(\Delta_1 + \Delta_2)^{1/2} + (\Delta_1 + 4\Delta_2)^{1/2}] \frac{\beta}{(\cos Q)^{1/2}} \end{aligned} \quad (27)$$

where

$$\begin{aligned} \Delta_1 &= (\eta_1 - 1)(\eta_2 - \cos Q) \cos Q \\ \Delta_2 &= \frac{1}{4}(\eta_2 - \cos Q \eta_1)^2 \end{aligned} \quad (28)$$

Straightforward calculations lead to the following low-temperature asymptotics of the chemical potentials η_1 and η_2

$$\begin{aligned} \eta_1 - 1 &= \frac{e^{-M_2\beta}}{2SJ(\cos Q)^{1/2}\beta} \left[\frac{3 + e^{-(M_1-M_2)\beta}}{9 - e^{-(M_1-M_2)\beta}} - \left(\frac{1 - e^{-(M_1-M_2)\beta}}{9 - e^{-(M_1-M_2)\beta}} \right)^{1/2} \right] \\ \eta_2 - \cos Q &= \frac{(\cos Q)^{1/2} e^{-M_2\beta}}{2SJ\beta} \left[\frac{3 + e^{-(M_1-M_2)\beta}}{9 - e^{-(M_1-M_2)\beta}} + \left(\frac{1 - e^{-(M_1-M_2)\beta}}{9 - e^{-(M_1-M_2)\beta}} \right)^{1/2} \right] \end{aligned} \quad (29)$$

where

$$M_1 = 4\pi S J m_1; \quad M_2 = 4\pi S J m_2 \cos Q \quad (30)$$

and m_1, m_2 are given by

$$\begin{aligned} 2m_1 &= 2S + 1 - \delta - (\cos Q)^{1/2} \\ 2m_2 &= 2S + 1 - \delta - (\cos Q)^{-1/2} \end{aligned} \quad (31)$$

For $Q = 2\pi$, $m_1 = m_2 = m$, $M_1 = M_2$, $\eta_1 = \eta_2 = \eta$ and

$$\eta - 1 = \frac{1}{4SJ\beta} e^{-4\pi S m} \quad (32)$$

For the other values of Q , $M_2 < M_1$. Q_2 is defined from Eq. (31) with $m_2 = 0$

$$Q_2 = \arccos \frac{1}{(2S + 1 - \delta)^2} \quad (33)$$

Finally, let Q runs the interval $Q_1 < Q < Q_2$. Then $\eta_1(T)$ goes to one, when T goes to zero, and $\eta_2(O) > |\cos Q|$.

$$\eta_1 - 1 = \frac{1}{(4SJ\beta)^2 (\eta_2 - \cos Q)} e^{-4\pi S J m_1 \beta} \quad (34)$$

where m_1 and $\eta_2(O)$ are determined by the equations

$$\begin{aligned} 2S + 1 - \delta &= \frac{1}{N} \sum_k \left(\frac{\eta_2 + \cos Q \gamma_k}{1 + \gamma_k} \right)^{1/2} + 2m_1 \\ 2S + 1 - \delta &= \frac{1}{N} \sum_k \left(\frac{1 + \gamma_k}{\eta_2 + \cos Q \gamma_k} \right)^{1/2} \end{aligned} \quad (35)$$

and $m_2 = 0$.

Let us recapitulate the results we have received above. When the temperature goes to zero $\eta_1(T)$ goes to one for all values of Q . At the same time $\eta_2(O) = |\cos Q|$ if $\pi \leq Q \leq Q_1$, and m_1, m_2 can be found from Eqs.(26). When $Q_1 < Q < Q_2$, $m_2 = 0$ and $\eta_2(O)$ and m_1 can be obtained from the system Eqs.(35). Finally, when $Q_2 \leq Q \leq 2\pi$, $\eta_2 = \cos Q$ and m_1, m_2 are given by Eqs. (31).

For the further convenience we have used a common notation for the variables m_1, m_2 when $\pi \leq Q \leq Q_1$ and when $Q_2 \leq Q \leq 2\pi$, but their origin is different. At zero temperature, when $\pi \leq Q \leq Q_1$, $\eta_1 = 1$ and $\eta_2 = |\cos Q|$, the bose system condenses at the wave vectors $k^* = (\pm\pi, \pm\pi)$ and $k_o = (0, 0)$. Then, the spin-wave correlators read

$$\begin{aligned} \langle a_k^+ a_k \rangle &= \frac{1}{2} \left[\frac{\epsilon_b(k)}{E_b(k)} \text{cth} \frac{\beta}{2} E_b(k) - 1 \right] (1 - \delta_{k,k^*}) (1 - \delta_{k,k_o}) \\ &\quad + \frac{1}{2} n_{k^*} \delta_{k,k^*} + \frac{1}{2} n_{k_o} \delta_{k,k_o} \end{aligned} \quad (36)$$

$$\begin{aligned} \langle a_k a_{-k} \rangle &= \langle a_k^+ a_{-k}^+ \rangle = \\ &= -\frac{D_k}{E_b(k)} \text{cth} \frac{\beta}{2} E_b(k) (1 - \delta_{k,k^*}) (1 - \delta_{k,k_o}) - \frac{1}{2} n_{k^*} \delta_{k,k^*} + \frac{1}{2} n_{k_o} \delta_{k,k_o} \end{aligned}$$

and

$$m_1 = \frac{1}{N} \sum_{k^*} n_{k^*}, \quad m_2 = \frac{n_{k_o}}{N} \quad (37)$$

When $Q_2 \leq Q \leq 2\pi$, $\eta_1 = 1$ and $\eta_2 = \cos Q$ the bose system condenses at the wave-vector k^* and for the spin-wave correlators are obtains

$$\begin{aligned} \langle a_k^+ a_k \rangle &= \frac{1}{2} \left[\frac{\epsilon_b(k)}{E_b(k)} \text{cth} \frac{\beta}{2} E_b(k) - 1 \right] (1 - \delta_{k,k^*}) + \frac{1}{2} n_{k^*}^{(1)} \delta_{k,k^*} \\ \langle a_k a_{-k} \rangle &= \langle a_k^+ a_{-k}^+ \rangle = -\frac{D_k}{E_b(k)} \text{cth} \frac{\beta}{2} E_b(k) (1 - \delta_{k,k^*}) - \frac{1}{2} n_{k^*}^{(2)} \delta_{k,k^*} \end{aligned} \quad (38)$$

where

$$\begin{aligned} \frac{1}{2N} \sum_{k^*} (n_{k^*}^{(1)} + n_{k^*}^{(2)}) &= m_1 \\ \frac{1}{2N} \sum_{k^*} (n_{k^*}^{(1)} - n_{k^*}^{(2)}) &= m_2 \end{aligned} \quad (39)$$

When $Q_1 < Q < Q_2$ the spin-wave correlators are given by Eq.(38) with $n_{k^*}^{(1)} = n_{k^*}^{(2)}$ ($m_2 = 0$).

To complete our analysis we shall discuss the magnon spectrum as a function of Q . From Eqs (23) we obtain that when $\pi \leq Q \leq Q_1$ there are

two antiferromagnetic magnons: $E_b(k) \simeq v_s |k - k^*|$ when $|k - k^*| \rightarrow 0$ and $E_b(k) \simeq v_s |k - k^o|$ when $|k - k^o| \rightarrow 0$, with spin-wave velocity $v_s = 4SJ\sqrt{|\cos Q|/2}$. When Q exceeds Q_1 one of the magnons opens a gap $E_b \simeq v_s\sqrt{\Delta^2 + |k - k^o|^2}$ where $\Delta^2 = 4(\eta_2 - |\cos Q|)/|\cos Q|$. In the interval $(3\pi/2, Q_2)$ there is only one magnon $E_b \simeq v'_s |k - k^*|$ with $v'_s = 2SJ(\eta_2 - \cos Q)^{1/2}$. The spin-wave velocity v'_s approaches zero when Q approaches Q_2 and in the interval $[Q_2, 2\pi]$ we obtain the ferromagnetic magnon $E_b \simeq SJ\sqrt{\cos Q} |k - k^*|^2$.

The angle Q is a free parameter in the theory. To determine it we shall calculate the energy of the system per site.

From Eqs.(5), taking into account the conditions Eqs.(12,14), one obtains in spin-wave approximation

$$\begin{aligned} & \langle \vec{S}_i \cdot \vec{S}_j \rangle = \\ & \cos(\theta_i - \theta_j) \left[\langle a_i^+ a_j \rangle \langle a_j^+ a_i \rangle + \langle a_i^+ a_j^+ \rangle \langle a_i a_j \rangle - \frac{1}{4} \langle \psi_i^+ \psi_j \rangle \langle \psi_j^+ \psi_i \rangle \right] \end{aligned} \quad (40)$$

$$\begin{aligned} & \langle c_{i\sigma}^+ c_{j\sigma} + h.c. \rangle = \\ & = -\cos \frac{\theta_i - \theta_j}{2} \left[\left(S + \frac{1}{2} \delta \right) (\langle \psi_j^+ \psi_i \rangle + \langle \psi_i^+ \psi_j \rangle) + \right. \\ & \quad \left. + \langle \psi_j^+ \psi_i \rangle \langle a_i^+ a_j \rangle + \langle \psi_i^+ \psi_j \rangle \langle a_j^+ a_i \rangle \right] \end{aligned} \quad (41)$$

and

$$\langle n_i n_j \rangle = (2S - \delta)^2 - \langle \psi_i^+ \psi_j \rangle \langle \psi_j^+ \psi_i \rangle$$

Putting Eqs.(40-42) into Eq.(1) one obtains for the energy per site at $T = 0$ and $\pi \leq Q \leq Q_2$

$$\begin{aligned} E = & -\frac{J}{2}(2S - \delta)^2 + \frac{1}{2}(1 - \cos Q) \left[\frac{1}{N} \sum_k \gamma_k \theta(-\epsilon_f(k)) \right]^2 + \\ & + \frac{J \cos Q}{4} \left[2S + 1 - \delta - \frac{1}{N} \sum_k (1 - \gamma_k) \left(\frac{1 + \gamma_k}{\eta_2 + \cos Q \gamma_k} \right)^{1/2} \right]^2 + \end{aligned}$$

$$\begin{aligned}
& + \frac{J \cos Q}{4} \left[2S + 1 - \delta - \frac{1}{N} \sum_k (1 + \gamma_k) \left(\frac{\eta_2 + \cos Q \gamma_k}{1 + \gamma_k} \right)^{1/2} \right]^2 + \\
& + 2t \cos \frac{Q}{2} \left[\frac{1}{N} \sum_k \gamma_k \theta(-\epsilon_f(k)) \right] \left[\frac{1}{2N} \sum_k (1 + \gamma_k) \left(\frac{\eta_2 + \cos Q \gamma_k}{1 + \gamma_k} \right)^{1/2} - \right. \\
& \quad \left. - \frac{1}{2N} \sum_k (1 - \gamma_k) \left(\frac{1 + \gamma_k}{\eta_2 + \cos Q \gamma_k} \right)^{1/2} + 2S + \delta \right] \quad (42)
\end{aligned}$$

where η_2 is defined above and $\theta(-\epsilon_f)$ is the step function.

When $Q_2 \leq Q \leq 2\pi$, the energy is given by

$$\begin{aligned}
E = & -\frac{J}{2}(2S - \delta)^2 + \frac{1}{2}(1 - \cos Q) \left[\frac{1}{N} \sum_k \gamma_k \theta(-\epsilon_f(k)) \right]^2 \\
& + \frac{J \cos Q}{4} \left[2S + 1 - \delta - \frac{1}{\sqrt{\cos Q}} \right]^2 + \\
& + \frac{J \cos Q}{4} \left[2S + 1 - \delta - \sqrt{\cos Q} \right]^2 + \\
& + 2t \cos \frac{Q}{2} \left[\frac{1}{N} \sum_k \gamma_k \theta(-\epsilon_f(k)) \right] \left[2\delta - 1 + \frac{1}{2\sqrt{\cos Q}} + \frac{\sqrt{\cos Q}}{2} \right]
\end{aligned}$$

For a given δ E is a function of Q , and the physical value of $Q(Q_o)$ is that which minimizes the energy. It depends on δ .

When the doping is very small Q_o is close to π , $\eta_2 = |\cos Q|$, and

$$\frac{1}{N} \sum_k \gamma_k \theta(-\epsilon_f(k)) \simeq \delta$$

The minimum of the energy is at

$$Q_o = \pi + \frac{S}{(S + 0.079)^2} \frac{t}{J} \delta \quad (43)$$

The energy per site is plotted in Fig.1 as a function of Q , for different values of δ , $t/J = 5$ and $S = 1/2$. When δ increases the minimum of the energy moves smoothly. A second minimum at $Q = 2\pi$ appears when the doping is large enough. This minimum decreases when δ increases and at some value of the doping the both minimums become equal. At this point Q_o

has a step-wise behavior. The curves $Q_o(\delta)$ are plotted in Fig. 2 for different values of t/J and $S = 1/2$. We treat the appearance of the second minimum as a appearance of instability and as a upper bound of applicability of our approach. For $t/J = 10$ this bound is $\delta = 0.31$, for $t/J = 5$; $\delta = 0.39$, for $t/J = 3$; $\delta = 0.47$ and for $t/J = 0, 5$; $\delta = 0.6$.

III. The spin-spin correlator

Let us consider the spin-spin correlation function in the limit of large separation between the two sites. At zero temperature the bose system condenses. Then, the large distance asymptotic of the bose correlators is determined by the values of their Fourier transforms at the wave vectors at which the system condenses. From Eqs.(36-39) one obtains

$$\langle a_i^+ a_j \rangle \langle a_j^+ a_i \rangle + \langle a_i a_j \rangle \langle a_i^+ a_j^+ \rangle \simeq m^2 \quad (44)$$

where

$$m^2 = \frac{1}{2}(m_1^2 + m_2^2) \quad (45)$$

The fermi term goes to zero when $|r_i - r_j|$ goes to infinity and we shall drop it. As a result we obtain that the large distance asymptotic of spin-spin correlator at $T = 0$ is

$$\langle \vec{S}_i \cdot \vec{S}_j \rangle = m^2 \cos Q_o \cdot (r_i - r_j) \quad (46)$$

This implies the presence of the long-range spiral phase. From Eqs.(26) one obtains that in leading order of δ

$$m_1 = m_2 = m = S - 0.197 - \frac{1}{2}\delta \quad (47)$$

In Fig. 3 we have plotted m as a function of δ for different values of t/J .

At finite temperature, the bose energy $E_b(k)$ (see Eq. (23)) has a finite gap. As a consequence, the bose correlators have a finite correlation length. For the spin-spin correlation function at small doping (when $\pi \leq Q_o(\delta) \leq$

$Q_2(\delta)$ and large distance one obtains

$$\begin{aligned} \langle \vec{S}_i \cdot \vec{S}_j \rangle \simeq & \frac{1}{(2SJ_\beta)^2} \frac{\cos Q_o \cdot (r_i - r_j)}{2\pi |r_i - r_j|} \left[\xi_1 e^{-\frac{|r_i - r_j|}{\xi_1}} + \right. \\ & \left. + \frac{\xi_2}{(\cos Q_o)^2} e^{-\frac{|r_i - r_j|}{\xi_2}} \right] \end{aligned} \quad (48)$$

where

$$\xi_1^{-1} = 4\sqrt{\eta_1 - 1} \quad (49)$$

$$\xi_2^{-1} = 4\sqrt{(\eta_2 - |\cos Q_o|)/|\cos Q_o|} \quad (50)$$

From Eqs. (25) it follows that ξ_1 and ξ_2 grow exponentially when T goes to zero. At very small doping $\eta_1 \simeq \eta_2$, $\xi_1 \simeq \xi_2$, $|\cos Q_o| \simeq 1$ and the spin-spin correlator looks like in the theory of the Heisenberg antiferromagnet⁸. When δ increases m_2 decreases and the first term in Eq.(48) is dominated when T approaches zero.

IV. Concluding remarks

In conclusion, we have investigated the two-dimensional $t - J$ model by modified spin-wave theory. A spin-spin correlator, we have obtained, at $T = 0$ Eq. (46) shows a spiral long-range order. At $T \neq 0$ the correlation function decays exponentially as in the theory of the Heisenberg antiferromagnet. The results, at small doping, are in qualitative agreement with the results obtained by Schwinger boson slave-fermion mean field theory⁵, except for the vector $3/2$ known from the earlier publications. Substantial quantitative discrepancies appear when δ increases. In spite of this the approach we have discussed above seems to be appropriate for the investigation of the normal state of the novel superconductors.

As a by-product we obtain the spin-stiffness constant ρ_S of the square lattice Heisenberg antiferromagnets. It can be defined by

$$\rho_S = \left. \frac{d^2 E}{dQ^2} \right|_{Q=\pi}$$

where $E(Q)$ is the energy per site of the system. From Eq. (42) at $\delta = 0$ one obtains

$$\rho_S = 2J(S + 0.079)^2$$

References

1. P.W.Anderson, Science, 235, 1196 (1987).
2. C.Gross, R.Joynt, and T.M.Rice, Phys.Rev. B36, 381 (1987).
3. F.Zhang and T.M.Rice, Phys.Rev. B37, 3759 (1988).
4. D.Arovas and A.Auerbach, Phys.Rev. B38, 316 (1988);
D.Yoshioka, J.Phys.Soc.Jpn. 58, 32 (1989).
5. D.Yoshioka, J.Phys.Soc.Jpn. 58, 1516 (1989);
C.Jayaprahash, H.R.Krishnamurty, and S.Sarker, Phys.Rev. B40, 2610 (1989);
K.Feisenberg, P.Hedegård, and M.Pederson, Phys.Rev. B40, 850 (1989).
6. B.Shraiman and E.Siggia, Phys.Rev.Lett 62, 1564 (1989).
7. M.Takahashi, Prog.Theor.Phys. Suppl. 87, 233 (1986); Phys.Rev.Lett. 58, 168 (1987).
8. M.Takahashi, Phys.Rev. B40, 2494 (1989).
9. S.B.Chakravarty, B.I.Halperin, and D.Nelson, Phys.Rev. B39, 2344 (1989).
10. N.I.Karchev, Theor. and Math. Phys. October 1992.
11. A.Angellucci and R.Link, MIT preprint, CTP# 2043 (1991).
12. J.E.Hirsh and S.Tang, Phys.Rev. B40, 4769 (1989).

FIGURE CAPTIONS

Fig. 1. The energy of the system per site as a function of Q for $t/J = 5$, $S = 1/2$ and different values of doping.

Fig. 2. The physical value Q_o of the spirality angle as a function of doping, for $t/J = 10$ (solid line with circles), $t/J = 5$ (dashed line), $t/J = 3$ (dotted line), $t/J = 0.5$ (dot-dashed line) and $S = 1/2$.

Fig. 3. The quantity m as a function of doping for $t/J = 10$ (solid line), $t/J = 5$ (dashed line), $t/J = 3$ (dotted line), $t/J = 0.5$ (dot-dashed line) and $S = 1/2$.